Langer-Giedion syndrome

Langer-Giedion syndrome is a condition that causes bone abnormalities and distinctive facial features. People with this condition have multiple noncancerous (benign) bone tumors called osteochondromas. Multiple osteochondromas may result in pain, limited range of joint movement, and pressure on nerves, blood vessels, the spinal cord, and tissues surrounding the osteochondromas. Affected individuals also have short stature and cone-shaped ends of the long bones (epiphyses). The characteristic appearance of individuals with Langer-Giedion syndrome includes sparse scalp hair, a rounded nose, a long flat area between the nose and the upper lip (philtrum), and a thin upper lip. Some people with this condition have loose skin in childhood, which typically resolves with age. Affected individuals may have some intellectual disability.

Frequency

Langer-Giedion syndrome is a rare condition; its incidence is unknown.

Genetic Changes

Langer-Giedion syndrome is caused by the deletion or mutation of at least two genes on chromosome 8. Researchers have determined that the loss of a functional *EXT1* gene is responsible for the multiple osteochondromas seen in people with Langer-Giedion syndrome. Loss of a functional *TRPS1* gene may cause the other bone and facial abnormalities. The *EXT1* gene and the *TRPS1* gene are always missing or mutated in affected individuals, but other neighboring genes may also be involved. The loss of additional genes from this region of chromosome 8 likely contributes to the varied features of this condition.

Langer-Giedion syndrome is often described as a contiguous gene deletion syndrome because it results from the loss of several neighboring genes.

Inheritance Pattern

Most cases of Langer-Giedion syndrome are not inherited, but occur as random events during the formation of reproductive cells (eggs or sperm) in a parent of an affected individual. These cases occur in people with no history of the disorder in their family. There have been very few instances in which people with Langer-Giedion syndrome have inherited the chromosomal deletion from a parent with the condition.

Langer-Giedion syndrome is considered an autosomal dominant condition because one copy of the altered chromosome 8 in each cell is sufficient to cause the disorder.

Other Names for This Condition

- Giedion-Langer Syndrome
- tricho-rhino-phalangeal syndrome type II
- trichorhinophalangeal syndrome type II
- TRPS II

Diagnosis & Management

Genetic Testing

 Genetic Testing Registry: Langer-Giedion syndrome https://www.ncbi.nlm.nih.gov/gtr/conditions/C0023003/

General Information from MedlinePlus

- Diagnostic Tests https://medlineplus.gov/diagnostictests.html
- Drug Therapy https://medlineplus.gov/drugtherapy.html
- Genetic Counseling https://medlineplus.gov/geneticcounseling.html
- Palliative Care https://medlineplus.gov/palliativecare.html
- Surgery and Rehabilitation https://medlineplus.gov/surgeryandrehabilitation.html

Additional Information & Resources

MedlinePlus

 Health Topic: Bone Diseases https://medlineplus.gov/bonediseases.html

Genetic and Rare Diseases Information Center

 Trichorhinophalangeal syndrome type 2 https://rarediseases.info.nih.gov/diseases/7801/trichorhinophalangeal-syndrometype-2

Educational Resources

- MalaCards: trichorhinophalangeal syndrome, type ii http://www.malacards.org/card/trichorhinophalangeal_syndrome_type_ii_2
- Orphanet: Langer-Giedion syndrome http://www.orpha.net/consor/cgi-bin/OC_Exp.php?Lng=EN&Expert=502
- Univeristy of Kansas Medical Center Resource List http://www.kumc.edu/gec/support/langer_g.html

Patient Support and Advocacy Resources

- Chromosome Disorder Outreach http://chromodisorder.org/
- National Organization for Rare Disorders (NORD)
 https://rarediseases.org/rare-diseases/trichorhinophalangeal-syndrome-type-ii/
- The MAGIC Foundation https://www.magicfoundation.org/
- Unique: Rare Chromosome Disorder Support Group http://www.rarechromo.org/html/home.asp

ClinicalTrials.gov

ClinicalTrials.gov
 https://clinicaltrials.gov/ct2/results?cond=%22Langer-Giedion+syndrome%22

Scientific Articles on PubMed

PubMed

https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28Langer-Giedion+syndrome %5BTIAB%5D%29+OR+%28tricho-rhino-phalangeal+syndrome+type+II%5BTIAB %5D%29+OR+%28trichorhinophalangeal+syndrome+type+II%5BTIAB%5D %29%29+OR+%28TRPS+II%5BTIAB%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D

OMIM

 TRICHORHINOPHALANGEAL SYNDROME, TYPE II http://omim.org/entry/150230

Sources for This Summary

- Hilton MJ, Sawyer JM, Gutiérrez L, Hogart A, Kung TC, Wells DE. Analysis of novel and recurrent mutations responsible for the tricho-rhino-phalangeal syndromes. J Hum Genet. 2002;47(3):103-6.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/11950061
- Riedl S, Giedion A, Schweitzer K, Müllner-Eidenböck A, Grill F, Frisch H, Lüdecke HJ. Pronounced short stature in a girl with tricho-rhino-phalangeal syndrome II (TRPS II, Langer-Giedion syndrome) and growth hormone deficiency. Am J Med Genet A. 2004 Dec 1;131(2):200-3.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/15523607
- Shanske AL, Patel A, Saukam S, Levy B, Lüdecke HJ. Clinical and molecular characterization of a patient with Langer-Giedion syndrome and mosaic del(8)(q22.3q24.13). Am J Med Genet A. 2008 Dec 15;146A(24):3211-6. doi: 10.1002/ajmg.a.32615.
 Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/19012352

Reprinted from Genetics Home Reference:

https://ghr.nlm.nih.gov/condition/langer-giedion-syndrome

Reviewed: February 2009 Published: March 21, 2017

Lister Hill National Center for Biomedical Communications U.S. National Library of Medicine National Institutes of Health Department of Health & Human Services